# Proposal

1. Research Title: (ภาษาอังกฤษ) Developing protocols for rationing scarce critical-care resources (intensive care unit (ICU) beds and ventilators) during the COVID-19 pandemic in Thailand (ภาษาไทย) การพัฒนาแนวทางแบ่งปันทรัพยากรทางการแพทย์ที่ขาดแคลนในช่วงการระบาด ของโควิด 19 ในประเทศไทย

## 2. Research members

#### 2.1 List of Researchers

1.	Dr. Yot Teerawattananon	Researcher
2.	Dr. Wanrudee Isaranuwatchai	Research assistant
3.	Ms. Rachel Archer	Research assistant
4.	Ms. Kanchanok Sirison	Research assistant
5.	Ms. Aniqa Islam Marshall	Research assistant
6.	Mr. Somtanuek Chotchoungchatchai	Research assistant
7.	Ms. Woranan Witthayapipopsakul	Research assistant
8.	Mr. Pisit Sriakkpokin	Research assistant

Address: Health Intervention and Technology Assessment Program 6 fl., 6 bldg.,

Department of Health, Ministry of Public Health, Tiwanon rd., Muang, Nonthaburi 11000,

Thailand

# 2.2 Project advisor

1.Dr. Viroj Tangcharoensathin Consultant

Address: International Health Policy Program (IHPP), Ministry of Public Health, Tiwanon rd., Muang, Nonthaburi 11000, Thailand

# 3. Introduction

# Current situation of COVID-19

The emerging pandemic outbreak caused by a newly discovered coronavirus (COVID-19) is considered a significant threat to the global public health (1). The first cases of COVID-19 can be traced back to Wuhan, China, in December 2019 (2). COVID-19 then spread rapidly to other Chinese provinces as well as neighboring countries and has since reached all but a few corners of the world (3, 4). The number of confirmed cases surpassed 1 million cases on 2nd April 2020 (5). Patients with COVID-19 may be asymptomatic or exhibit mild to severe symptoms, with an estimated fatality rate of approximately 2%

among the medically attended patients (6). Currently, there are no specific drugs or vaccine available for treating this disease.

In Thailand, the number of confirmed COVID-19 cases is rising. As of 2<sup>nd</sup> April 2020, there have been 1,875 COVID-19 confirmed cases of which 23 severe cases are currently receiving intensive care treatment (5). For mild cases of COVID-19, initiating respiratory support is neither needed nor recommended. Once the disease progresses, pneumonia, severe pneumonia, acute respiratory distress syndrome (ARDS), and sepsis could be observed within the first few weeks of onset, requiring prompt respiratory support and intensive care to increase chance of recovery (7). The clinical management of patients with COVID-19 is considered to be similar to other viral infections such as Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS), as well as sepsis and organ failure (8, 9). The initiating respiratory support is recommended to alleviate interstitial pneumonia characterized by severe hypoxia (10). Increasing number of hospitalized COVID-19 patients places a higher demand of clinical resources including intensive critical units (ICU) and mechanical ventilators.

Bangkok has experienced the highest number of COVID-19 cases in Thailand. A report by the Department of Medical Services showed that as of 8PM on 3<sup>rd</sup> April 2020, there were 109 Airborne Infection Isolation Rooms (AIIR)-ICU beds across 90 hospitals in Bangkok and surrounding areas. At the same time, 101 AIIR-ICU beds were occupied and 8 were vacant. Based on the forecasting model, as of 2<sup>nd</sup> April 2020, it is estimated that the demand for ICU beds in the capital will reach 190 beds by 19<sup>th</sup> April 2020 and by the end of April, it could reach over 300 beds, exceeding the supply of available AIIR-ICU beds in the greater Bangkok area. Whilst plans are in place for the creation of more AIIR-ICU beds, it is considered inadequate for the increasing demand (7). For ventilators' status, it is reported that 10,184 ventilators have been reserved throughout the country (7).

## Current rationing process

Whilst the COVID-19 outbreak in Thailand remains a rapidly evolving situation, the current trajectory predicts a surge in patients requiring critical care (severe COVID-19 cases) which will result in a shortage of ICU beds and ventilators. The COVID-19 global pandemic has shown that even the most well-resourced health systems experience shortages of critical care resources in the height of the outbreak. In Lombardy, Italy, for instance, there were not enough ventilators to go around despite large efforts to increase supply (11). In preparation for the exceptional circumstances where demand for critical care resources will outstrip supply, there is an urgent need to establish explicit protocols to determine how resources are to be rationed and reallocated (12). This process, known as triage, will determine the

priority of access to critical care in the current pandemic (13). In these special circumstances where ICU beds and ventilators are scarce, the practice of rationing is unavoidable. The question therefore is not whether one should ration or not, rather how to ration in a fair and explicit manner or do so 'in the heat of the moment' (14).

There is a wealth of literature stating that bedside clinicians should not be tasked with deciding who gets access to live-saving care (12, 14, 15). Placing the responsibility of triage on the clinicians who are treating the patients can be morally distressing and mentally debilitating and is likely to result in varied criteria across hospitals which is ethically problematic (14). It is recommended that protocols on rationing be issued by the provincial/national level authorities and be uniformly applied across health facilities (14). This can make sure that rationing decisions are informed by wider societal values (e.g. protecting vulnerable populations, maximizing health benefit etc.) and capture resource-allocation trade-offs (11, 14). Furthermore, for an effective triage process, protocols need to be concise and easy-to-understand by health professionals, patients and their caregivers (16). Health professionals, technical experts (including ethicists) and the general public all need to be engaged in the development of the criteria and mechanism for rationing to ensure it reflects the priorities of society and to maintain public trust in the health system process. The protocols should not be developed in a 'closed room' manner (17). Given the current COVID-19 pandemic, this research is timely and will assist hospitals in implementing a fair and transparent approach to allocating scarce critical care resources that are high in-demand and are the last resource for saving lives of these severely affected patients.

# 4. Objectives

The main objective is to develop mechanisms for rationing both ICU beds and ventilators and establish criteria for allocation of these scarce resources that can be applied fairly and consistently across Thailand. These protocols (criteria and mechanisms) will be applicable at the point of demand exceeding supply in the exceptional circumstances brought about by the COVID-19 pandemic in Thailand.

The target audience for this work range from clinicians and Ministry of Public Health (MOPH) officials to patients, caregivers and the general public. The protocols will be developed in consultation with all relevant stakeholders to support the response efforts of the MOPH. The work will be disseminated to all stakeholders for wider use. The expected output is clear and concise protocols that are informed by a wealth of literature, guidelines, expert opinion and the values of the Thai society.

## 5. Research methods

#### A) Literature review

A literature review will be carried out using academic databases including PubMed and Science Direct, as well as ethical recommendations and existing international guidelines from countries such as Germany, Japan, South Korea, Switzerland, United Kingdom, United States of America. The project team will simultaneously review the critical-care triage policy (for ICU beds and ventilators) and guidance on resource allocation and bed flow management in Thailand.

#### B) Interviews with health professionals

HITAP will conduct qualitative interviews with the healthcare professionals who are working in ICU and non-ICU units in selected hospitals - Ramathibodi Hospital, Siriraj Hospital, Rajavithi Hospital, and Petcharavej Hospital. Given the current outbreak, the sample of interviewees will be kept minimal (e.g. 3-5 doctors and nurses) as the study should not disrupt current services. A semi-structured interview questionnaire will be developed which will adhere to the ethics standards in Thailand and will be pretested to ensure the instrument is valid and captures all the relevant aspects.

## C) Develop the protocols and publish online

C1) Following activities (A) and (B), the protocol on the rationing ICU beds and ventilators in response to the COVID-19 outbreak will be developed by the research team under the supervision and guidance of the steering committee chaired by Dr.Viroj Tangcharoensathin. The first draft protocol will be consulted with stakeholders consisting of public health specialists, ethicists, psychologists, political scientists, medical lawyer, medical specialists, medical anthropologists and religious leaders.

C2) The second draft protocol will be developed following the consultation with the stakeholders. Once the second draft has been created, another stakeholders consultation will be arranged for the protocol finalization process. For protocol evaluation, the draft protocol will be published online for general public.

# D) Present findings to Ministry of Public Health

The research team will convene a meeting with the MOPH to present a collective summary of the findings in response to the research objectives. The meeting will provide an opportunity to elicit expert input on the criteria and mechanism for critical-care resource allocation.

# E) Final report

A final report will be prepared to describe the work that was undertaken and document the project findings. The report will contain the criteria and mechanism for allocation critical care resources (specifically ICU beds and ventilators) under exceptional conditions raised by the COVID-19 pandemic in Thailand and perhaps future outbreaks.

## F) Manuscript in a peer-reviewed journal

A manuscript will be submitted and published in an open-access peer-reviewed journal on the critical care resource allocation protocols developed in response to the COVID-19 pandemic in Thailand. The manuscript will summarize lessons learned from this exercise and prospects for the future.

## 6. Benefits from this research

- 1. This study will support health care professionals, hospitals and policymakers for the allocation of critical-care scarce resources under a public health emergency.
- 2. This study will help relieve a moral burden upon attending physicians and ensure that the decision is fair, transparent, ethically appropriate and consistent throughout.

#### Inclusion criteria

- Participants agree and are willing to be interviewed.
- Participants sign an informed consent form.
- Participants are ICU and non-ICU doctors and nurses from public and private hospitals.

## Exclusion criteria

• Participants withdraw consent from the interview.

## 7. <u>Timeline</u>

The estimated timeline for the tasks A-D is six weeks from project inception.

## 8. Ethical issues

There is minimal risk about personal information of interviewees. The research team will store data on a computer with an access code. The researchers will explain the details of the research project including its objectives, main approaches and expected benefits to the interviewees every time prior to signing the informed consent forms. The researchers are aware that the interviewees have the right to

accept or decline to participate in this research. In addition, in case the interviewees would like to withdraw their consent during the interview, the researchers will discontinue the interview immediately.

All personal information, sound records and interview transcripts of the interviewees will be kept confidential. The study results will be reported in form of overall narrative description, which cannot be tracked back to the individual informants.

# 12. ประวัติและประสบการณ์ของหัวหน้าโครงการ คณะผู้วิจัย และที่ปรึกษา

d	ตำแหน่ง	หน่วยงานที่สังกัด พร้อมที่อยู่ เบอร์ โทรศัพท์	อีเมล	ความรับผิดชอบต่อโครงการที่ เสนอ			
ชื่อ-นามสกุล				บทบาทหน้าที่	คิดเป็น สัดส่วน <b>)</b> %)	FTE	คุณวุฒิ และสาขาที่เชี่ยวชาญ
ดร. นพ.ยศ ตีระวัฒนานนท์	หัวหน้า โครงการ	โครงการประเมิน เทคโนโลยีและนโยบาย ด้านสุขภาพ 02-590-4549	yot.t@hitap.net	พัฒนาข้อเสนอ โครงการวิจัย ดำเนินการวิจัย และ วิเคราะห์ข้อมูล	30	0.50	การประเมินเทคโนโลยีและนโยบายด้าน สุขภาพ Doctor of Philosophy in Health Economics, University of East Anglia, England, 2006
Ms. Rachel Archer	ผู้วิจัยร่วม	โครงการประเมิน เทคโนโลยีและนโยบาย ด้านสุขภาพ 02-590-4549	rachel.a@hitap.net	ประสานงาน ดำเนินการวิจัยและ วิเคราะห์ข้อมูล	10	0.50	การประเมินเทคโนโลยีและนโยบายด้าน สุขภาพ University of Sheffield   Masters of Public Health & International Development

น.ส.กานต์ชนก ศิริสอน	ผู้วิจัยร่วม	โครงการประเมิน เทคโนโลยีและนโยบาย ด้านสุขภาพ 02-590- 4549	kanchanok.s@hitap.net	ประสานงาน ดำเนินการวิจัยและ วิเคราะห์ข้อมูล	10	0.50	การประเมินเทคโนโลยีและนโยบายด้าน สุขภาพ Master of Pharmacy (MPharm), Durham University, UK
Ms. Aniqa Islam Marshall	ผู้วิจัยร่วม	สำนักงานพัฒนา นโยบายสุขภาพระหว่าง ประเทศ 02-02-590-2366-7	aniqa@ihpp.thaigov.net	ประสานงาน ดำเนินการวิจัยและ วิเคราะห์ข้อมูล	10	0.50	พัฒนานโยบายสุขภาพระหว่างประเทศ  Master of Science in Public Health (MSPH) Johns Hopkins Bloomberg School of Public Health
นายสมธนึก โชติช่วงฉัตรชัย	ผู้วิจัยร่วม	สำนักงานพัฒนา นโยบายสุขภาพระหว่าง ประเทศ 02-02-590-2366-7	Somtanuek@ihpp.thaigov.net , tueyz@hotmail.com	ประสานงาน ดำเนินการวิจัยและ วิเคราะห์ข้อมูล	10	0.10	พัฒนานโยบายสุขภาพระหว่างประเทศ  Doctor of Philosophy Program in Phytopharmaceutical Sciences, Faculty of Pharmacy, Mahidol University, Thailand, 2012-present, Topic: An efficacy of honey and Thai herbal oil preparation on pressure ulcer treatment using hybrid methodology
น.ส.วรณัน วิทยาพิภพสกุล	ผู้วิจัยร่วม	สำนักงานพัฒนา นโยบายสุขภาพระหว่าง ประเทศ	woranan88@gmail.com, woranan@ihpp.thaigov.net	ประสานงาน ดำเนินการวิจัยและ วิเคราะห์ข้อมูล	10	0.10	พัฒนานโยบายสุขภาพระหว่างประเทศ

		02-590-2366-7					MSc Health Systems and Public Policy (Merit class)
นายพิสิษฐ์ ศรีอัคคโภคิน		สำนักงานคณะกรรมการ สุขภาพแห่งชาติ	large_urchin@hotmail.com	ประสานงาน ดำเนินการวิจัยและ วิเคราะห์ข้อมูล	10	0.10	ผู้ชำนาญการ ทำงานกฎหมายการแพทย์ใน เรื่องสิทธิและหน้าที่ด้านสุขภาพ ตาม พระราชบัญญัติสุขภาพแห่งชาติ พ.ศ. 2550 ปริญญาโทสาธารณสุขศาสตร์ มหาวิทยาลัยมหิดล
คร. นพ.วิโรจน์ ตั้งเจริญเสถียร	ที่ปรึกษา	สำนักงานพัฒนา นโยบายสุขภาพระหว่าง ประเทศ 02-590-2366-7	Viroj@ihpp.thaigov.net	ดำเนินการวิจัย และ วิเคราะห์ข้อมูล	10	0.10	พัฒนานโยบายสุขภาพระหว่างประเทศ London School of Hygiene and Tropical Medicine, University of London

## <u>References</u>

- 1. Pneumonia of unknown cause China: disease outbreak news. Geneva: World Health Organization, January 5, 2020. Available from https://www.who.int/csr/don/05-january-2020-pneumonia-of-unkown-cause-china/en/.
- 2. Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus–infected pneumonia. N Engl J Med. 2020; 382: 1199-1207.
- 3. Vincent JM, Marion K, van Doremalen N, et al. A novel coronavirus emerging in China—key questions for impact assessment. N Engl J Med. 2020.
- 4. She J, Jiang J, Ye L, et al. Clin Trans Med 2019 novel coronavirus of pneumonia in Wuhan, China: emerging attack and management strategies. 2020; 9: 19.
- 5. EOC-DDC Thailand. COVID-19 (EOC-DDC Thailand) live report, April 3, 2020. Available from: https://ddcportal.ddc.moph.go.th/portal/apps/opsdashboard/index.html#/20f3466e075e45e5946aa87c96e 8ad65 [Accessed 3rd April 2020].
- 6. Lipsitch M, Donnelly CA, Fraser C, et al. Potential Biases in Estimating Absolute and Relative Case-Fatality Risks during Outbreaks. PLoS Negl Trop Dis 2015;9(7): e0003846-e0003846.
- 7. Sirilak S. An estimation of Covid-19 report in Thailand. MOPH Intelligence Unit (MIU). Published: April 3, 2020
- 8. Alhazzani, W., Møller, M.H., Arabi, Y.M. et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). Intensive Care Med (2020).
- 9. Poston JT, Patel BK, Davis AM. Management of Critically Ill Adults With COVID-19. JAMA. Published online March 26, 2020.
- 10. Cascella M, Rajnik M, Cuomo A, et al. Features, Evaluation and Treatment Coronavirus (COVID-19) [Updated 2020 Mar 20]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-.
- 11. Rosenbaum, Facing Covid-19 in Italy Ethics, Logistics, and Therapeutics on the Epidemic's Front Line, The New England Journal of Medicine, March 18, 2020, DOI:10.1056/NEJMp2005492
- 12. White DB, Lo B. A Framework for Rationing Ventilators and Critical Care Beds During the COVID-19 Pandemic. JAMA. March 27, 2020. doi:10.1001/jama.2020.5046

- 13. Ghanbari et al., Ethical prioritization of patients during disaster triager: A systematic review of current evidence, International Emergency Nursing, 2019, DOI: 10.1016/j.ienj.2018.10.004
- 14. Emanuel et al., Fair Allocation of Scarce Medical Resources in the Time of Covid-19, The New England Journal of Medicine, March, 23 2020.
- 15. Truog et al., The Toughest Triage Allocating Ventilators in a Pandemic, The New England Journal of Medicine, March 23, 2020, DOI:10.1056/NEJMp2005689
- 16. Baker et al., Essential care of critical illness must not be forgotten in the COVID-19 pandemic. The Lancet. April 1, 2020, DOI: 10.1016/S0140-6736(20)30793-5
- 17. Alicia Ely Yamin and Ole F. Norheim, 3 Human Rights Imperatives for Rationing Care in the Time of Coronavirus, Petrie-Flom Blog, March 27, 2020.